

The AGN Source Population in the Chandra Deep Field-North Survey: Constraints from X-ray Spectroscopy and Variability

F. E. BAUER,¹ C. VIGNALI,¹ D. M. ALEXANDER,¹ W. N. BRANDT,¹ G. P. GARMIRE,¹
A. E. HORNSCHMEIER,¹ P. BROOS,¹ L. TOWNSLEY,¹ AND D. P. SCHNEIDER¹

¹ Department of Astronomy & Astrophysics, 525 Davey Lab, The Pennsylvania State University, University Park, PA 16802.

1. Introduction

With the 0.5–10.0 keV background nearly resolved (e.g., Giacomini et al. 2000; Cowie et al. 2002), the emphasis has shifted towards understanding the nature of the faint X-ray population — a clear necessity if we wish to understand the formation and evolution of active galactic nuclei (AGN). Here we report on preliminary spectral and temporal X-ray studies of AGN in the 2 Ms Chandra Deep Field-North Survey (CDF-N; D.M. Alexander et al. in prep). Our sample is comprised of 136 CDF-N sources with > 200 net counts ($\sim 30\%$ of the total CDF-N sample), spanning $(0.014\text{--}2) \times 10^{-13}$ ergs $\text{cm}^{-2} \text{s}^{-1}$; 72 sources have known redshifts.

2. Spectra

We initially fitted each source with a fixed Galactic N_{H} ($1.6 \times 10^{20} \text{ cm}^{-2}$; Stark et al. 1992), a variable intrinsic N_{H} at the source redshift if known, and a simple powerlaw with variable photon index Γ . Approximately 70% of the sources fitted by this model have $\chi^2_{\nu} < 1.3$, with the remaining sources often exhibiting soft excess residuals (i.e., due to a partial covering or thermal component). The median spectral parameters are $\Gamma = 1.61$ and $N_{\text{H}} = 1.7 \times 10^{21} \text{ cm}^{-2}$ for all 136 sources, or $\Gamma = 1.55$ and $N_{\text{H}} = 2.6 \times 10^{21} \text{ cm}^{-2}$ for the 61 sources with > 500 counts. The intrinsic N_{H} distribution for the sample and a comparison of L_{X} vs. N_{H} are shown in Fig. 1. The N_{H} distribution is strongly skewed towards low/Galactic values with only a small fraction of sources lying above 10^{23} cm^{-2} . The 27 optically identified broad-line AGN (BLAGN) trace the overall sample.

3. Emission Lines

Ten of the 136 AGN ($\approx 7\%$) exhibit obvious Fe K α emission-line features, with equivalent widths (EWs) of 0.1–1.3 keV. Two of the emission-line sources appear to be Compton thick AGN, displaying both large EWs and extremely flat spectral slopes $\Gamma < 1.0$ characteristic of pure reflection (e.g., Maiolino et al. 1998). We have also constrained the number of potential Compton thick sources among the 72 sources with known redshifts. By adding a Gaussian component in the




Fig. 1. Left: The overall intrinsic N_{H} distribution for our sample. Right: L_{X} vs. N_{H} for our sample. Sources without known redshifts are plotted as L_{X} lower limits assuming $z = 0.5$.

source spectrum ($E = 6.4 \text{ keV}$, $\sigma = 0 \text{ keV}$), we can place upper limits on the Fe K α EWs. Only 8 of the 72 sources ($\approx 11\%$) have 90% confidence EW upper limits above 1 keV, and, of these, none has a measured photon index $\Gamma < 1.0$. Since $\sim 50\%$ of all known Compton thick sources have $\Gamma < 1.0$ and large EWs can also arise from anisotropic ionizing radiation or variability lags between continuum and line emissions (e.g., Maiolino et al. 1998; Bassani et al. 1999), the true number of Compton thick sources among our spectroscopically identified sources may be even smaller.

4. Variability

The 20 individual CDF-N observations span approximately 27 months and offer an unprecedented probe of long-term X-ray variability in distant AGN. Using both Kolmogorov-Smirnov and χ^2 statistics, we find that $\sim 94\%$ of the BLAGN, $\sim 90\%$ of the > 500 count sources, and $\sim 60\%$ of the overall sample show indications of variability.

Acknowledgments

We gratefully acknowledge the financial support of NSF CAREER award AST-9983783 (FEB, CV, DMA, WNB), NASA grant NAS 8-38252 (GPG, PI), NASA GSRP grant NGT5-50247 (AEH), and NSF grant AST-9900703 (DPS).

References

- Bassani, L. et al. 1999, ApJS, 121, 473
- Cowie, L.L. et al. 2002, ApJL, 566, 5
- Giacconi, R. et al. 2001, ApJ, 551, 624
- Maiolino, R. et al. 1998, A&A, 338, 781
- Stark, A. et al. 1992, ApJS, 79, 77